

## Residents of States with Swine Influenza Cases

CDC has identified human cases of swine influenza A (H1N1) virus infection in people in the U.S. CDC is working with local and state health agencies to investigate these cases. We have determined that this virus is contagious and is spreading from human to human. However, at this time, we have not determined how easily the virus spreads between people. As with any infectious disease, we are recommending precautionary measures for people residing in these areas.

- Cover your nose and mouth with a tissue when you cough or sneeze. Throw the tissue in the trash after you use it.
- Wash your hands often with soap and water, especially after you cough or sneeze. Alcohol-based hands cleaners are also effective.
- Try to avoid close contact with sick people.
- If you get sick, CDC recommends that you stay home from work or school and limit contact with others to keep from infecting them.
- Avoid touching your eyes, nose or mouth. Germs spread that way.

There is no vaccine available at this time, so it is important for people living in these areas to take steps to prevent spreading the virus to others. If people are ill, they should attempt to stay at home and limit contact with others. Healthy residents living in these areas should take [everyday preventive actions](#).

People who live in these areas who develop an illness with fever and respiratory symptoms, such as cough and runny nose, and possibly other symptoms, such as body aches, nausea, or vomiting or diarrhea, should contact their health care provider. Their health care provider will determine whether influenza testing is needed.

## Clinicians

Clinicians should consider the possibility of swine influenza virus infections in patients presenting with febrile respiratory illness who

1. Live in an area where human cases of swine influenza A (H1N1) has been identified or
2. Have traveled to an area where human cases of swine influenza A (H1N1) has been identified or
3. Have been in contact with ill persons from these areas in the 7 days prior to their illness onset.

If swine flu is suspected, clinicians should obtain a respiratory swab for swine influenza testing and place it in a refrigerator (not a freezer). Once collected, the clinician should contact their state or local health department to facilitate transport and timely diagnosis at a state public health laboratory.

Source: <http://www.cdc.gov/flu/>